

INSTRUCTIONS FOR USE

ZELSVUMI™ (zel-SOOV-mee) (berdazimer) topical gel, 10.3%




(berdazimer) topical gel, 10.3%

161545-2


This Instructions for Use contains information on how to prepare and apply ZELSVUMI. Read this Instructions for Use before you start using ZELSVUMI and each time you get a new prescription. Apply ZELSVUMI exactly as your healthcare provider tells you to. Ask your healthcare provider if you have any questions.

Box Contents (Figure A):


1 Tube A



1 Tube B



1 Dosing Guide



Important Information You Need to Know Before Applying ZELSVUMI:

- ZELSVUMI is for use on top of the skin (for topical use only).**
 - Do not treat bumps close to your eye or get gel in your eye. If gel gets in your eye, rinse with water.
 - Do not eat gel. If gel is swallowed, contact your healthcare provider.
 - Do not apply gel to open wounds. If gel gets into an open wound, rinse with water.
 - Do not get gel in your mouth, vagina, or on areas of your skin where you do not have bumps.
- ZELSVUMI comes in a box that contains (See Figure A):
 - 1 blue tube that contains berdazimer gel (Tube A)
 - 1 yellow tube that contains hydrogel (Tube B)
 - 1 Dosing Guide
- If you have trouble treating the bumps as the instructions describe, contact your healthcare provider.
- The gels in Tube A and Tube B must be mixed together on the Dosing Guide before applying to your skin.**
- Do not mix the gels until you are ready to apply ZELSVUMI.**
- Make sure to put the blue cap back onto the blue tube and put the yellow cap onto the yellow tube. If you mix the caps, wipe caps with **dry** tissue and replace caps on correct tubes.
- Hold tubes gently. If excess gel leaks out, wipe with tissue or cloth.

- Apply mixed gel to clean, dry skin.
- After mixing the gel, apply it to your skin right away.
- Dosing Guide**
 - Only mix gels on Dosing Guide** to ensure the gels are measured accurately and mixed thoroughly. **Do not mix gels directly on skin or in palm of hand.**
 - Always lay Dosing Guide on a flat surface.
 - Always use the Dosing Guide to dispense gels.
 - Throw away any ZELSVUMI remaining on the Dosing Guide after use. **Do not store mixed ZELSVUMI.**
 - Do not** throw away the Dosing Guide or Instructions for Use. You will re-use the Dosing Guide and Instructions for Use for every application.
 - If your Dosing Guide, Instructions for Use, or tubes are lost or damaged, please contact your healthcare provider, pharmacy, or manufacturer at 1-855-330-7546.

Preparing to Apply ZELSVUMI

- On a clean, flat surface, open the ZELSVUMI box and remove Tube A and Tube B.
- Break the seal on tubes. You only need to do this the first time you open the tubes.
 - Take cap off Tube A and break seal by pushing the pointed end of the cap into the foil.
 - Put cap back on Tube A tightly.
 - Repeat the above steps with Tube B.

Turn to the back page for more instructions.

Instructions begin on the other side.

Applying ZELSVUMI

Step 1. Wash Hands

Always wash and dry hands before applying gel.



Step 2. Dispense Gel


Place Dosing Guide on a flat surface.

Remove cap from Tube A. Hold Tube A near blue Tube A dosing lane.

Squeeze gel onto blue lane to cover **entire area of the lane**. The gel is white on the Dosing Guide. Screw cap back onto Tube A.

Remove cap from Tube B. Hold Tube B near yellow Tube B dosing lane.

Squeeze gel onto yellow lane to cover **entire area of the lane**. The gel is clear to almost clear on the Dosing Guide. Screw cap back onto Tube B.



Step 3. Mix Gels

Use a fingertip to combine the 2 gels from the blue and yellow lanes. Mix the gels in the center of the Dosing Guide using a circular motion. It is important that the 2 gels get mixed well. Mix the 2 gels together while slowly counting to 20. You may see clumps during mixing and this is normal. Apply the mixed gel right away to the bumps.




Step 4. Apply Mixed Gel

Apply an even thin layer of mixed gel to each bump right away.

- All bumps should be treated with gel.

Make sure to cover bumps that are new, hard to reach, or out of sight.



If you run out of mixed gel before you have treated all the bumps, wipe off your Dosing Guide with a **dry** tissue. Follow steps 2, 3, and 4 to apply the mixed gel to the remaining untreated bumps.

After all bumps have been treated:

- Follow the cleaning and storing steps below.
- Wait at least 10 minutes after applying gel before putting clothes on skin to allow ZELSVUMI to dry.
- Wait at least 1 hour after applying gel before swimming, washing, bathing, or showering.

Step 5. Clean Dosing Guide

Use water and mild soap to remove remaining gel from Dosing Guide. Dry Dosing Guide.



Step 6. Storing ZELSVUMI

Put dry Dosing Guide and this Instructions for Use back in box. Secure caps on tubes and place back in box.

Store ZELSVUMI at room temperature between 68°F to 77°F (20°C to 25°C) in a dry location. Product contains alcohol and should be kept away from open flame. Do not freeze. Throw away if not used within 60 days after receiving ZELSVUMI.

Step 7. Wash Hands

Always wash hands after you place the contents back in the box unless your hands were treated.



Disposing of ZELSVUMI

Throw away (dispose of) tubes in household trash when empty or when treatment is stopped.



Manufactured for:

LNHC, Inc., Durham, NC 27703

 **Zelsuvmi**™
(berdazimer) topical gel, 10.3%